

KAON CONDENSATION IN DENSE STELLAR MATTER*Chang-Hwan Lee^a and Mannque Rho^b

- a) *Department of Physics and Center for Theoretical Physics
Seoul National University, Seoul 151-742, Korea*
- b) *Service de Physique Théorique, CEA Saclay
91191 Gif-sur-Yvette Cedex, France*

ABSTRACT

This talk is based on work done in collaboration with G.E. Brown and D.-P. Min on kaon condensation in dense baryonic medium treated in chiral perturbation theory using heavy-baryon formalism. It contains, in addition to what was recently published, some new results based on the analysis on kaonic atoms by Friedman, Gal and Batty and a discussion on a renormalization-group analysis to meson condensation made together with H.K. Lee and Sin. Negatively charged kaons are predicted to condense at the critical density $2 \lesssim \rho/\rho_0 \lesssim 4$, in the range to allow all the intriguing new phenomena predicted by Brown and Bethe to take place in compact star matter.

*Based on talk given by MR at *International Symposium on "Strangeness and Quark Matter"*, September 1-5, 1994, Krete, Greece and at *YITP Workshop "From Hadronic Matter to Quark Matter: Evolving View of Hadronic Matter"*, October 30-November 1, 1994, Yukawa Institute, Kyoto University, Kyoto, Japan.

1 Motivation

Recent work by Bethe and Brown[1] on the maximum mass of stable compact stars – called “neutron stars” in the past but more appropriately “nuclear (or nucleon) stars” – suggest that the nuclear equation of state (EOS) in the interior of compact stars must be considerably softened at densities a few times the nuclear matter density ρ_0 by one or several hadronic phase transitions. It is now fairly clear that neither pion condensation nor quark matter will figure at a density low enough to be relevant to the star matter although the issue is not yet completely settled. As Bethe and Brown suggest, kaon condensation could however take place at a density 3–4 times the normal matter density and hence play an important role in explaining the remarkably narrow range of compact star masses observed in nature[2].

The aim of this talk is to describe an honest calculation that predicts the critical density for kaon condensation. The strategy is to take up what Kaplan and Nelson[3] started, namely chiral perturbation theory (χ PT). Kaplan and Nelson predicted in tree order of χ PT that kaons condense in neutron matter at $\rho \lesssim 3\rho_0$. Our calculation goes to next-to-next-to-leading (NNL) order. It turns out that the calculation confirms the Kaplan-Nelson prediction although in the process new and interesting physical elements are uncovered. Our result is that for reasonable ranges of parameters involved, the critical density comes out to be

$$2 \lesssim \rho/\rho_0 \lesssim 4. \tag{1}$$

This is the range of densities relevant to the Bethe-Brown scenario for the formation of light-mass black holes for stars that exceed the critical mass of $M = 1.5M_\odot$.

2 Where Do the Kaons Come From?

There are two situations where the production of kaons brings out interesting physics. One is their properties in relativistic heavy-ion collisions that involve temperature. Here kaon condensation is not directly relevant but the *mechanism* that triggers kaon condensation in the relevant situation has intriguing consequences on the properties of kaons observed in heavy-ion experiments. This is discussed in a recent review[4] and will not be discussed here. What we are interested in is what kaons do in cold dense matter appropriate to compact objects that result from the collapse of massive stars.

In stellar collapse, as matter density ρ increases, the electron chemical potential μ_e (determined by the chemical potentials of neutrons and protons in the system together with charge neutrality) increases, reaching several hundreds of MeV. If the electron chemical potential reaches the “effective mass” of a meson Φ , m_Φ , then the electron can “decay” into

a Φ as[5]

$$e^- \rightarrow \Phi^- + \nu_e. \quad (2)$$

In nature, the only low-mass bosons are the pseudo-Goldstone bosons $\Phi^- = \pi^-, K^-$. While lowest in mass, the pions do not seem to play an important role, so the next possible boson is the kaon with its mass ~ 500 MeV in free space. The electron chemical potential cannot reach this high, so on-shell kaons cannot be produced by this process. However as will be described below, the kaon in medium can undergo a mass shift due to density-dependent renormalization. As the μ_e increases and the effective kaon mass M_K^* decreases as ρ increases, the process (2) can occur at some density ρ_c . Kaons so produced will Bose-condense at that density ρ_c . Whether or not this will occur then depends on whether or not M_K^* will decrease enough in density so that it meets μ_e . Such a condensation will be of physical interest if the critical density is low enough and the energy gain is high enough. This is the possibility we shall address below.

3 Baryon Chiral Perturbation Theory

3.1 *Effective Field Theory for Nuclear Matter*

The process we are interested in requires a field theory that can describe simultaneously normal nuclear matter and phase transitions therefrom. The most relevant ingredient of QCD that is needed here is spontaneously broken chiral symmetry. We are specifically interested in chiral $SU(3) \times SU(3)$ symmetry since strangeness is involved. In order to address the problem, we need to start from a realistic effective chiral Lagrangian, obtain a nuclear matter of the right properties from it and then determine whether strangeness condensation occurs.

Unfortunately we do not yet know how to describe nuclear matter starting from a chiral Lagrangian. There are various suggestions and one promising one is that nuclear matter arises as a solitonic matter from a chiral effective action, a sort of chiral liquid[6] resembling Landau Fermi liquid. The hope is that the resulting effective action would look like Walecka's mean-field model. There is as yet no convincing derivation along this line. In the work reported here, we will have to assume that we have a nuclear matter that comes out of an effective chiral action. Given such a ground state containing no strange degrees of freedom, we would like to study fluctuations along the strangeness direction and determine if instability along that direction develops signaling a phase transition. We are therefore assuming that we can get the properties of normal nuclear matter from phenomenology, that is, that nuclear matter is a Fermi-liquid fixed point[7, 8]. In principle, a precise knowledge of this ground state from a chiral effective Lagrangian at a nonperturbative QCD level would

allow us to determine the coefficients that appear in the effective Lagrangian with which to describe fluctuations around the soliton background – i.e., the Fermi liquid – and with which we could then compute all nuclear response functions. At present such a derivation does not exist. In a recent paper by Brown and one of the authors (BR91)[10], it is assumed that in medium at a matter density $\rho \sim \rho_0$, the *nuclear* effective field theory can be written in terms of the medium-dependent coupling constants g^* and masses of hadrons m^* while preserving the free-space structure of a sigma model. This leads to the so-called Brown-Rho scaling. In BR91[10], the nonlinear sigma model implemented with trace anomaly of QCD is used to arrive at the scaling law. The precise way that this scaling makes sense is elaborated by Adami and Brown[9] and in the review (BR94)[4]. There have been numerous papers written with some of the essential points of this scaling misinterpreted.

Given such an effective field theory, we can make a general argument on the stability in various flavor directions of nuclear matter at high density. This can be done along the line of arguments developed for condensed matter physics by Shankar[7] and Polchinski[8] using renormalization group flow. We sketch the essential argument following Lee, Rho and Sin[11].

What we are interested in is whether the system in question develops instability along the direction of strangeness and if so, by which physical mechanism. This analysis will not give us the critical density. The critical density will be calculated by using chiral perturbation theory. For this purpose we will focus on the kaon frequency near the electron chemical potential. By Baym's theorem[12], one can identify the kaon chemical potential associated with charge conservation, μ_K , with the electron chemical potential, μ_e , which we shall simply write μ in what follows. This means that we will be looking at the vicinity of $\omega \sim \mu$ in the kaon dispersion formula. We shall assume that

$$|\omega - \mu| < \mu. \quad (3)$$

As mentioned, we assume that nucleons in nuclear matter are in Fermi-liquid state with the Fermi energy μ_F and the Fermi momentum k_F . Define ψ as the nucleon field fluctuating around the Fermi surface such that the momentum integral has a cut-off Λ_N ,

$$k_F - \Lambda_N < |\vec{k}| < k_F + \Lambda_N. \quad (4)$$

Kaons can interact with the nucleons through three-point functions of the KNN type (Yukawa interaction) and through four-point interactions of the $KKNN$ type. We shall consider s-wave kaon-nucleon interactions, for which the Yukawa interaction can be ignored. A generic action involving the nucleon field ψ and the kaon field Φ can then be written, schematically, as

$$S = \int d\omega d^3q \Phi^*(\omega, \vec{q}) \left(\omega - q^2/2\mu_K \right) \Phi(\omega, \vec{q}) - \int d\omega d^3q \tilde{M}_K \Phi^* \Phi$$

$$\begin{aligned}
& + \int (d\omega d^3q)^2 (d\epsilon d^3k)^2 h \Phi^* \Phi \psi^\dagger \psi \delta^4(\omega, \epsilon, \vec{q}, \vec{k}) \\
& + \int d\epsilon d^3k \psi^\dagger (\epsilon - \epsilon(k)) \psi + g \int (d\epsilon d^3k)^4 \psi^\dagger \psi^\dagger \psi \psi \delta^4(\epsilon, \vec{k})
\end{aligned} \tag{5}$$

where $\tilde{M} = (M_K^2 - \mu^2)/2\mu$ and h and g are constants. The four-Fermi interaction with the coefficient g stands for Fermi-liquid interactions in nuclear matter. (In nuclear matter, one can have four such terms because of the nucleon spin and isospin degrees of freedom. We need not specify them for our purpose.) This is a toy action but it is generic in that the results of χ PT we will obtain below can be put into this form.

The renormalization group flow of this action can be analyzed in the following way. Since we are assuming that nuclear matter is a Fermi-liquid fixed point, fluctuations in the non-strange direction in the nucleon sector are stable: The four-Fermi interaction g is irrelevant or at best marginal. Fluctuations in the strange direction involve the kaon field Φ . Suppose we have integrated out all the high-frequency modes above the cut-off Λ measured with respect to μ . We are interested in the stability of the system under the renormalization group transformation $\Lambda \rightarrow s\Lambda$ ($s < 1$) as $s \rightarrow 0$. A scaling analysis shows that the interaction term h is irrelevant while the “mass term” $\tilde{M}$ is relevant. The renormalization group-flow of the “mass term” and the interaction term h can be readily written down and solved[11] (with $t = -\ln s$),

$$\tilde{M}(t) = (\tilde{M}_0 - \frac{Dh_0}{1+a})e^t + \frac{Dh_0}{1+a}e^{-at} \tag{6}$$

with

$$h(t) = h_0 e^{-at}, \quad h_0 \geq 0 \tag{7}$$

where $D = \frac{3(1+\alpha^2)\alpha}{2\mu}\rho_N > 0$, $\alpha = \Lambda/k_F > 0$ and $a = 1/2$. We see from Eq.(6) that as $s \rightarrow 0$ for which $h \rightarrow 0$, $\tilde{M}$ changes sign for some $(\tilde{M}_0, h_0 \geq 0)$. Thus although irrelevant, an attractive interaction h_0 determines the direction of the mass flow whereas it is the “mass term” that drives the system to instability.

3.2 Chiral Counting

Armed with the general information on the instability in the strangeness direction, we now calculate the critical density in χ PT. Remember that we are to look at the instability in the kaon direction, so it suffices for us to look at the fluctuations around the Fermi-liquid state. For this we need an effective chiral Lagrangian involving baryons as well as Goldstone bosons. When baryons are present, χ PT is not as firmly formulated as when they are absent[13]. The reason is that the baryon mass m_B is $\sim \Lambda_\chi \sim 1$ GeV, the chiral symmetry

breaking scale. It is more expedient, therefore, to redefine the baryon field so as to remove the mass from the baryon propagator

$$B_v = e^{im_B \gamma \cdot v} P_+ B \quad (8)$$

where $P_+ = (1 + \gamma \cdot v)/2$ and write the baryon four-momentum

$$p_\mu = m_B v_\mu + k_\mu \quad (9)$$

where k_μ is the small residual momentum indicating the baryon being slightly off-shell. When acted on by a derivative, the baryon field B_v yields a term of $O(k)$. Chiral perturbation theory in terms of B_v and Goldstone bosons ($\pi \cdot \lambda/2$) is known as “heavy-baryon (HB) χ PT” [14]. HB χ PT consists of making chiral expansion in derivatives on Goldstone boson fields, ∂_M/Λ_χ , and on baryon fields, ∂_B/m_B , and in the quark mass matrix, $\kappa\mathcal{M}/\Lambda_\chi^2$. In the meson sector, this is just what Gasser and Leutwyler did for $\pi\pi$ scattering. In the baryon sector, consistency with this expansion requires that the chiral counting be made with $B^\dagger(\cdots)B$, not with $\bar{B}(\cdots)B$. This means that in medium, it is always the baryon density $\rho(r)$ that comes in and *not* the scalar density $\rho_s(r)$. This point seems to be misunderstood by some workers in the field, a cause for one of the red herrings in the literature.

Following Weinberg[15], we organize the chiral expansion in power Q^ν where Q is the characteristic energy/momentum scale we are looking at ($Q \ll \Lambda_\chi$) and

$$\nu = 4 - N_n - 2C + 2L + \sum_i \Delta_i \quad (10)$$

with the sum over i running over the vertices that appear in the graph and

$$\Delta_i = d_i + \frac{1}{2}n_i - 2. \quad (11)$$

Here ν gives the power of small momentum (or energy) for a process involving N_n nucleon lines, L number of loops, d_i number of derivatives (or powers of meson mass) in the i th vertex, n_i number of nucleon lines entering into i th vertex and C is the number of separate connected pieces of the Feynman graph. Chiral invariance requires that $\Delta_i \geq 0$, so that the leading power is given by $L = 0$, $\nu = 4 - N_n - 2C$.

As an example, consider KN scattering. The leading term here is the tree graph with $\nu = 1$ and with $N_n = C = 1$. The next order terms are $\nu = 2$ tree graphs with $\Delta = 1$ that involves two derivatives or one factor of the mass matrix $\mathcal{M}$. From $\nu = 3$ on, we have loop graphs contributing together with appropriate counter terms.

In considering kaon-nuclear interactions as in the case of kaon condensation, we need to consider the case with $N_n \geq 2$ and $C \geq 2$. In dealing with many-body system, one can simply fix $4 - N_n$ and consider C explicitly. For instance if one has two nucleons (for reasons

mentioned below, this is sufficient, with multinucleon interactions being suppressed), then we have $4 - N_n = 2$ but C can be 2 or 1, the former describing a kaon scattering on a single nucleon with a spectator nucleon propagating without interactions and the latter a kaon scattering irreducibly on a two-nucleon complex. Thus intrinsic n -nucleon processes are suppressed compared with $(n-1)$ -nucleon processes by at least $O(Q^2)$. This observation will be used later for arguing that four-Fermi interactions are negligible in kaon condensation. This is somewhat like the suppression of three-body nuclear forces[15] and of three-body exchange currents[16] in chiral Lagrangians.

3.3 Kaon-Nucleon Scattering

Given a chiral Lagrangian, we need to first determine the parameters of the Lagrangian from available phenomenology. This is inevitable in effective field theories. We shall first look at kaon-nucleon scattering at low energies. This was done by Lee *et al.*[17, 18] which we summarize here. We shall compute the scattering amplitude to one-loop order and this entails a Lagrangian written to $O(Q^3)$ as one can see from the Weinberg counting rule. Instead of writing it out in its full glory, we write it in a schematic form as

$$\mathcal{L} = \sum_i \mathcal{L}_i[B_\nu, U, \mathcal{M}] \quad (12)$$

where the subscript i stands for ν relevant to the KN channel. Here B_ν stands for both octet and decuplet baryons and U the Sugawara form for octet Goldstone bosons. For KN scattering in free-space, the Lagrangian is bilinear in the baryon field. Details are given in Lee *et al.* [17, 18]. Let us specify a few terms in (12) so as to streamline our discussion. Focusing on s-wave scattering, $\mathcal{L}_1$ contains the leading order term that may be described by the exchange of an ω between kaon and nucleon, attractive for K^-N and repulsive for K^+N and an isovector term corresponding to the exchange of a ρ meson. These terms are proportional to the kaon frequency ω . To next order, $\mathcal{L}_2$ contains the “ KN sigma term” proportional to Σ_{KN}/f^2 where f is the pion decay constant and a term proportional to ω^2 which may be saturated by decuplet intermediate states. The $\nu = 3$ pieces are counter terms that contain terms that remove divergences in the loop calculations and finite terms that are to be determined from experiments. The complete s-wave scattering amplitudes calculated to the NNL order come out to be

$$\begin{aligned} a_0^{K^\pm p} &= \frac{m_B}{4\pi f^2(m_B + M_K)} \left[\mp M_K + (\bar{d}_s + \bar{d}_v)M_K^2 + \{(L_s + L_v) \pm (\bar{g}_s + \bar{g}_v)\} M_K^3 \right] \\ &\quad + \delta a_{\Lambda^*}^{K^\pm p} \\ a_0^{K^\pm n} &= \frac{m_B}{4\pi f^2(m_B + M_K)} \left[\mp \frac{1}{2} M_K + (\bar{d}_s - \bar{d}_v)M_K^2 + \{(L_s - L_v) \pm (\bar{g}_s - \bar{g}_v)\} M_K^3 \right] \end{aligned} \quad (13)$$

where M_K is the kaon mass, m_B the baryon (nucleon) mass, $\bar{d}_s$ is the t-channel isoscalar contribution [$\mathcal{O}(Q^2)$], and $\bar{d}_v$ is the t-channel isovector one [$\mathcal{O}(Q^2)$], both coming from $\mathcal{L}_2$, $L_s(L_v)$ is the finite crossing-even t-channel isoscalar (isovector) finite one-loop contribution [$\mathcal{O}(Q^3)$] having the numerical values

$$L_s M_K \approx -0.109 \text{ fm}, \quad L_v M_K \approx +0.021 \text{ fm} \quad (14)$$

and the quantity $\bar{g}_s(\bar{g}_v)$ is the crossing-odd t-channel isoscalar (isovector) contribution [$\mathcal{O}(Q^3)$] from one-loop plus counter terms in $\mathcal{L}_3$. Finally $\delta a_{\Lambda^*}^{K^\pm p}$ is the contribution from the $\Lambda(1405)$ intermediate state Born diagram,

$$\delta a_{\Lambda^*}^{K^\pm p} = -\frac{m_B}{4\pi f^2(m_B + M_K)} \left[\frac{g_{\Lambda^*}^2 M_K^2}{m_B \mp M_K - m_{\Lambda^*}} \right] \quad (15)$$

which is completely determined given experimental data on the coupling g_{Λ^*} and the complex mass m_{Λ^*} .

There are four unknowns $\bar{d}_{s,v}$, $\bar{g}_{s,v}$ in (13) which can be determined from four experimental (real part of) scattering lengths

$$\begin{aligned} a_0^{K^+p} &= -0.31 \text{ fm}, & a_0^{K^-p} &= -0.67 + i0.63 \text{ fm} \\ a_0^{K^+n} &= -0.20 \text{ fm}, & a_0^{K^-n} &= +0.37 + i0.57 \text{ fm}. \end{aligned} \quad (16)$$

The results are

$$\begin{aligned} \bar{d}_s &\approx 0.201 \text{ fm}, & \bar{d}_v &\approx 0.013 \text{ fm}, \\ \bar{g}_s M_K &\approx 0.008 \text{ fm}, & \bar{g}_v M_K &\approx 0.002 \text{ fm}. \end{aligned} \quad (17)$$

So far, no prediction is made. However given the parameters so fixed, one can then go ahead and calculate the s-wave amplitude that enters in kaon condensation. This amounts to going off-shell in the ω variable, that is, in the kinematics where $\omega \neq M_K$. In doing this, one encounters an ambiguity due to the ω dependence of the coefficients $\bar{d}$ which consist of the “ KN sigma term” and “ ω^2 term” which get compounded on-shell into one term. In the calculation reported in Lee *et al.*[17, 18], we chose to fix the “ ω^2 term” by resonance saturation and leave the “sigma term” to be fixed by the on-shell data. The predicted off-shell amplitudes[17] agree reasonably with phenomenologically constructed off-shell amplitudes. All the constants of the chiral Lagrangian bilinear in the baryon field are thereby determined to $\mathcal{O}(Q^3)$.

3.4 Four-Fermi Interactions

In medium, the chiral Lagrangian can have multi-Fermi interactions as a result of “mode elimination.” Here we consider four-Fermi interactions, ignoring higher-body interactions. We shall see that this is justified.

As stated above, we need to focus on four-Fermi interactions that involve strangeness degrees of freedom. Nonstrange four-Fermi interactions are subsumed in the Fermi-liquid structure of normal nuclear matter. For s-wave kaon-nuclear interactions, we only have the $\Lambda(1405)$ to account for. There are only two terms,

$$\mathcal{L}_{4-fermion} = C_{\Lambda^*}^S \bar{\Lambda}_v^* \Lambda_v^* \text{Tr } \bar{B}_v B_v + C_{\Lambda^*}^T \bar{\Lambda}_v^* \sigma^k \Lambda_v^* \text{Tr } \bar{B}_v \sigma^k B_v \quad (18)$$

where $C_{\Lambda^*}^{S,T}$ are the dimension -2 (M^{-2}) parameters to be fixed empirically and σ^k acts on baryon spinor. We shall now describe how to fix these two parameters from kaonic atom data.

In order to confront kaonic atom data, we need to calculate the kaon self-energy Π in nuclei. The off-shell amplitude determined above gives the so-called “impulse” term

$$\Pi_K^{imp}(\omega) = - \left(\rho_p \mathcal{T}_{free}^{K^-p}(\omega) + \rho_n \mathcal{T}_{free}^{K^-n}(\omega) \right) \quad (19)$$

where $\mathcal{T}^{KN}$ is the off-shell s-wave KN transition matrix. (The amplitude $\mathcal{T}^{KN}$ taken on-shell, *i.e.*, $\omega = M_K$, and the scattering length a^{KN} are related by $a^{KN} = \frac{1}{4\pi(1+M_K/m_B)} \mathcal{T}^{KN}$.) Medium corrections to this “impulse” term, obtained from one-loop graphs by replacing the free-space nucleon propagator by the in-medium propagator, shall be denoted as

$$- \left(\rho_p \delta \mathcal{T}_{\rho_N}^{K^-p}(\omega) + \rho_n \delta \mathcal{T}_{\rho_N}^{K^-n}(\omega) \right). \quad (20)$$

These two terms (19) and (20) are completely determined by the parameters fixed above. The new parameters of the four-Fermi interaction come into play in the first two self-energy graphs of Fig.1 (the last two graphs do not involve four-Fermi interactions but enter at the same order; they are free of unknown parameters),

$$\begin{aligned} \Pi_{\Lambda^*}(\omega) = & -\frac{g_{\Lambda^*}^2}{f^2} \left(\frac{\omega}{\omega + m_B - m_{\Lambda^*}} \right)^2 \left\{ C_{\Lambda^*}^S \rho_p \left(\rho_n + \frac{1}{2} \rho_p \right) - \frac{3}{2} C_{\Lambda^*}^T \rho_p^2 \right\} \\ & + \frac{g_{\Lambda^*}^2}{f^4} \rho_p \left(\frac{\omega}{\omega + m_B - m_{\Lambda^*}} \right) \omega^2 \left\{ (2\Sigma_K^p(\omega) + \Sigma_K^n(\omega)) \right. \\ & \quad \left. - g_{\Lambda^*}^2 \left(\frac{\omega}{\omega + m_B - m_{\Lambda^*}} \right) (\Sigma_K^p(\omega) + \Sigma_K^n(\omega)) \right\} \quad (21) \end{aligned}$$

where g_{Λ^*} is the renormalized $KN\Lambda^*$ coupling constant determined in Lee *et al.*[17] and $\Sigma_K^N(\omega)$ is a known integral that depends on proton and neutron densities and M_K . Note that while the second term of (21) gives repulsion corresponding to a Pauli quenching, the first term can give either attraction or repulsion depending on the sign of $(C_{\Lambda^*}^S [\rho_n + \frac{1}{2} \rho_p] - \frac{3}{2} C_{\Lambda^*}^T \rho_p)$. For symmetric nuclear matter, only the combination $(C_{\Lambda^*}^S - C_{\Lambda^*}^T)$ enters in the self-energy. This is an important element for kaonic atom.




Figure 1: The in-medium two-loop kaon self-energy involving $\Lambda(1405)$. Figures a and b contain the constants of the four-Fermi interaction and figures c and d are Pauli corrections

The complete self-energy to in-medium two-loop order is then

$$\begin{aligned} \Pi_K(\omega) = & - \left(\rho_p \mathcal{T}_{free}^{K^-p}(\omega) + \rho_n \mathcal{T}_{free}^{K^-n}(\omega) \right) - \left(\rho_p \delta \mathcal{T}_{\rho_N}^{K^-p}(\omega) + \rho_n \delta \mathcal{T}_{\rho_N}^{K^-n}(\omega) \right) \\ & + \Pi_{\Lambda^*}(\omega). \end{aligned} \quad (22)$$

We now turn to fixing the constants of the four-Fermi interactions based on the recent analysis of kaonic atoms by Friedman, Gal and Batty [19]. For later purpose we shall parametrize the proton and neutron densities by the proton fraction x and the nucleon density $u = \rho/\rho_0$ as

$$\rho_p = x\rho, \quad \rho_n = (1-x)\rho, \quad \rho = u\rho_0. \quad (23)$$

Now Friedman *et al.*[19] found from their analysis that the optical potential for the K^- in medium has an attraction of the order of

$$\Delta V \equiv M_K^* - M_K \approx -(200 \pm 20) \text{ MeV at } u = 0.97 \quad (24)$$

with

$$M_K^* \equiv \sqrt{M_K^2 + \Pi_K}. \quad (25)$$

This implies approximately for $x = 1/2$

$$(C_{\Lambda^*}^S - C_{\Lambda^*}^T)f^2 \approx 20. \quad (26)$$

Friedman *et al.* [19] note that their “nominal” optical potential gives an attraction of order of 800 MeV when extrapolated to three times the normal density. We show in Table 1 what our theory predicts at higher densities than normal. At $u = 3$, the net attraction is only about 1.7 times the one at $u = 1$.

Table 1

K^- effective mass(M_K^*) and the attraction ($\Delta V \equiv M_K^* - M_K$) in symmetric nuclear matter ($x = 0.5$) as function of density u in unit of MeV for $(C_{\Lambda^*}^S - C_{\Lambda^*}^T)f^2 = 20$.

u	M_K^*	ΔV
0.5	348.6	-146.4
1.0	294.5	-200.5
1.5	249.3	-245.8
2.0	211.7	-283.3
2.5	179.9	-315.1
3.0	153.0	-342.0
3.5	129.0	-366.0
4.0	110.4	-384.6

Equation (21) shows that for symmetric nuclear matter ($x = 1/2$), the combination $(C_{\Lambda^*}^S + C_{\Lambda^*}^T)$ does not enter into the self-energy formula. In order to extract it as needed for non-symmetric system as in compact star matter, we need information for nuclei with $x \neq 1/2$. This can be done from the results of Friedman *et al.* by noting that our self-energy is nonlinear in x , so

$$\frac{\partial \Delta V}{\partial x}(C_{\Lambda^*}^S, \rho \approx \rho_0)|_{x=1/2} \approx 400 \, b_1/b_0 \, \text{MeV} \quad (27)$$

where $b_{0,1}$ are the constants given by Friedman *et al.*. This relation determines the coefficient $C_{\Lambda^*}^S$. The result is shown in Table 2 (first three columns).

Friedman *et al.*[19] find the acceptable value to be $b_1/b_0 = -0.56 \pm 0.82$. But there is one point which needs to be discussed in interpreting this number in the context of our theory. The constant $C_{\Lambda^*}^S$ shifts linearly the effective in-medium mass of $\Lambda(1405)$, with the mass shift being given by

$$\delta m_{\Lambda^*} = \sum_{i=a,b} \delta \Sigma_{\Lambda^*}^{(i)}(\omega = m_{\Lambda^*} - m_B) \quad (28)$$

where

$$\begin{aligned}\delta\Sigma_{\Lambda^*}^{(a)}(\omega) &= -\frac{g_{\Lambda^*}^2}{f^2}\omega^2 (\Sigma_K^p(\omega) + \Sigma_K^n(\omega)) \\ \delta\Sigma_{\Lambda^*}^{(b)}(\omega) &= -C_{\Lambda^*}^S(\rho_p + \rho_n).\end{aligned}\tag{29}$$

For nuclear matter density $u = 1$ and $x = 1/2$, the shift is

$$\delta m_{\Lambda^*}(u, x, y) \approx [62 - 150.3 \times y] \text{ MeV}\tag{30}$$

with $y = C_{\Lambda^*}^S f^2$. It seems highly unlikely that the $\Lambda(1405)$ will be shifted by hundreds of MeV in nuclear matter. This means that y must be of $O(1)$, and *not* $O(10)$. For $y = 0.41$ which corresponds to $b_1/b_0 \approx -0.4$, there is no shift at normal matter density. We believe this is a reasonable value. In fact, $y = 0$ is also acceptable. It would be interesting to measure the shift of $\Lambda(1405)$ to fix the constant $C_{\Lambda^*}^S$ more precisely although its precise magnitude seems to matter only a little for kaonic atoms and as it turns out, negligibly for kaon condensation.

Table 2

Determination of $C_{\Lambda^*}^S$ from the kaonic atom data[19] and the critical density (obtained with the constant so determined) for kaon condensation for various forms of symmetry energy $F(u)$. $y = 0.41$ corresponds to no $\Lambda(1405)$ mass shift in medium at the normal matter density.

$y = C_{\Lambda^*}^S f^2$	$\partial\Delta V/\partial x$	b_1/b_0	u_c		
			$F(u) = \frac{2u^2}{1+u}$	$F(u) = u$	$F(u) = \sqrt{u}$
50	125.78MeV	0.314	2.247	2.492	2.942
40	65.11MeV	0.163	2.320	2.572	3.051
30	4.44MeV	0.011	2.407	2.680	3.189
20	-56.23MeV	-0.141	2.528	2.821	3.372
10	-116.91MeV	-0.292	2.696	3.033	3.645
0.41	-175.089MeV	-0.438	2.958	3.391	4.159
0	-177.58MeV	-0.444	2.973	3.414	4.195
-10	-238.25MeV	-0.596	3.564	4.741	~ 5.897

Let us comment briefly on the role of multi-Fermion Lagrangians. This will eliminate another red herring in the literature. The Weinberg counting rule shows that the four-Fermi interactions are suppressed by $O(Q^2)$ relative to the terms involving bilinears of Fermi fields. In general n -Fermi interactions will be suppressed by the same order relative to $(n-1)$ -Fermi

interactions. In considering kaon condensation, what this means in conjunction with the renormalization-group flow argument, is that n -Fermi interactions with $n \geq 4$ are irrelevant in the RGE sense, and hence unimportant for condensation. The situation with the kaonic atom data is a bit different. While the strength of the four-Fermi interaction, y , is not important (this can be seen in Lee *et al.*[18], Table 3), its presence is essential for the attraction that seems to be required. This is in contrast to the kaon condensation which is driven by the “mass flow” with four-Fermi interactions being irrelevant in the RGE sense.

4 Kaon Condensation

We have now all the ingredients needed to calculate the critical density for negatively charged kaon condensation in dense nuclear star matter. For this, we will follow the procedure given in work of Thorsson, Prakash and Lattimer (TPL)[20]. As argued by Brown, Kubodera and Rho[5], we need not consider pions when electrons with high chemical potential can trigger condensation through the process $e^- \rightarrow K^- \nu_e$. Thus we can focus on the spatially uniform condensate

$$\langle K^- \rangle = v_K e^{-i\mu t}. \quad (31)$$

The energy density $\tilde{\epsilon}$ – which is related to the effective potential in the standard way – is given by,

$$\begin{aligned} \tilde{\epsilon}(u, x, \mu, v_K) = & \frac{3}{5} E_F^{(0)} u^{\frac{5}{3}} \rho_0 + V(u) + u \rho_0 (1 - 2x)^2 S(u) \\ & - [\mu^2 - M_K^2 - \Pi_K(\mu, u, x)] v_K^2 + \sum_{n \geq 2} a_n(\mu, u, x) v_K^n \\ & + \mu u \rho_0 x + \tilde{\epsilon}_e + \theta(|\mu| - m_\mu) \tilde{\epsilon}_\mu \end{aligned} \quad (32)$$

where $E_F^{(0)} = (p_F^{(0)})^2 / 2m_B$ and $p_F^{(0)} = (3\pi^2 \rho_0 / 2)^{\frac{1}{3}}$ are, respectively, Fermi energy and momentum at nuclear density. The $V(u)$ is a potential for symmetric nuclear matter as described by Prakash *et al.*[21] which is presumably subsumed in contact four-Fermi interactions (and one-pion-exchange – nonlocal – interaction) in the non-strange sector as mentioned above. It will affect the equation of state in the condensed phase but not the critical density, so we will drop it from now on. The nuclear symmetry energy $S(u)$ – also subsumed in four-Fermi interactions in the non-strange sector – does play a role as we know from Prakash *et al.*[21]: Protons enter to neutralize the charge of condensing K^- ’s making the resulting compact star “nuclear” rather than neutron star as one learns in standard astrophysics textbooks. We take the form advocated by Prakash *et al.*[21]

$$S(u) = \left(2^{\frac{2}{3}} - 1\right) \frac{3}{5} E_F^{(0)} \left(u^{\frac{2}{3}} - F(u)\right) + S_0 F(u) \quad (33)$$

where $F(u)$ is the potential contributions to the symmetry energy and $S_0 \simeq 30 \text{ MeV}$ is the bulk symmetry energy parameter. We use three different forms of $F(u)$ [21]

$$F(u) = u, \quad F(u) = \frac{2u^2}{1+u}, \quad F(u) = \sqrt{u}. \quad (34)$$

The contributions of the filled Fermi seas of electrons and muons are[20]

$$\begin{aligned} \tilde{\epsilon}_e &= -\frac{\mu^4}{12\pi^2} \\ \tilde{\epsilon}_\mu &= \epsilon_\mu - \mu\rho_\mu = \frac{m_\mu^4}{8\pi^2} \left((2t^2 + 1)t\sqrt{t^2 + 1} - \ln(t^2 + \sqrt{t^2 + 1}) \right) - \mu \frac{p_{F_\mu}^3}{3\pi^2} \end{aligned} \quad (35)$$

where $p_{F_\mu} = \sqrt{\mu^2 - m_\mu^2}$ is the Fermi momentum and $t = p_{F_\mu}/m_\mu$.

The ground-state energy prior to kaon condensation is then obtained by extremizing the energy density $\tilde{\epsilon}$ with respect to x , μ and v_K :

$$\left. \frac{\partial \epsilon}{\partial x} \right|_{v_K=0} = 0, \quad \left. \frac{\partial \epsilon}{\partial \mu} \right|_{v_K=0} = 0, \quad \left. \frac{\partial \epsilon}{\partial v_K^2} \right|_{v_K=0} = 0 \quad (36)$$

from which we obtain three equations corresponding, respectively, to beta equilibrium, charge neutrality and dispersion relation. The critical density so obtained is given for three different $F(u)$'s in Table 2. The result is

$$2 < u_c \lesssim 4. \quad (37)$$

5 Discussion

We note that the largest sensitivity is associated with the part that is not controlled by chiral symmetry, namely the density dependence of the symmetry energy function $F(u)$. This uncertainty reflects the part of interaction that is not directly given by chiral Lagrangians, that is, the part leading to normal nuclear matter. This is the major short-coming of our calculation.

Related to this issue is BR scaling. As we argued, were we able to derive nuclear matter from effective chiral Lagrangians, we would have parameters of the theory determined at that point reflecting the background around which fluctuations are to be made. The BR scaling was proposed in that spirit but with a rather strong assumption: That a sigma model governs dynamics in medium as in free space with only coupling constants and masses scaled a function of density. Up to date, no derivation of this scaling from basic principles has been made. In this sense, we might consider it as a conjecture. Suppose we apply BR scaling. The only way the procedure can make sense is to apply the scaling argument to the tree order terms, but not to the loop corrections. The result of this procedure is significant in

that the critical density is brought down in an intuitively plausible way to about $u_c \sim 2$, with very little dependence on parameters, loop corrections and multi-Fermi interactions. Thus slightly modified from (37), we arrive at the announced result

$$2 \lesssim u_c \lesssim 4. \quad (38)$$

Whether or not the Bethe-Brown scenario[1] of compact star formation is supported by the chiral Lagrangian approach will have to await the calculation of the equation of state at in-medium two-loop order, which is in progress. Our conjecture is that to the extent that our work confirms the original Kaplan-Nelson calculation[3], the compact star properties calculated previously at the tree level[20] would come out qualitatively unmodified in the higher-order chiral perturbation theory.

Acknowledgments

We would like to thank G.E. Brown, H.K. Lee, D.-P. Min and S.-J. Sin for discussions. The work of CHL is supported in part by the Korea Science and Engineering Foundation through the CTP of SNU and in part by the Korea Ministry of Education under Grant No. BSRI-94-2418.

References

- [1] G.E. Brown and H.A. Bethe, *Astrophys. J.* **423** (1994) 659; H.A. Bethe and G.E. Brown, "Observational constraints on the maximum neutron star mass," *Astrophys. J.*, to be published.
- [2] Z. Arzoumanian *et al.*, Aspen Winter Conf. on Astrophys., "Millisecond pulsars: A decade of surprises," PASP Conf. Ser. (1994); M.H. Van Kerkwijk, J. Van Paradijs, E.J. Zuidewijk, G. Hammerschlag-Hensberge, L. Laper and C. Sterken, submitted to *Astron. & Astrophys.* (1994).
- [3] D.B. Kaplan and A.E. Nelson, *Phys. Lett.* **B175** (1986) 57.
- [4] G.E. Brown and M. Rho, "Chiral restoration in hot and/or dense matter," *Phys. Repts.*, to appear.
- [5] G.E. Brown, K. Kubodera and M. Rho, *Phys. Lett.* **B192** (1987) 273.
- [6] B.W. Lynn, *Nucl. Phys.* **B402** (1993) 281.
- [7] R. Shankar, *Rev. Mod. Phys.* **66** (1994) 129.

- [8] J. Polchinski, in: *Recent Directions in Particle Physics*, eds. J. Harvey and J. Polchinski (World Scientific, Singapore, 1993) p. 235.
- [9] C. Adami and G.E. Brown, *Phys. Repts.* **224** (1993) 1.
- [10] G.E. Brown and M. Rho, *Phys. Rev. Lett.* **66** (1991) 2720.
- [11] H.K. Lee, M. Rho and S.-J. Sin, “Renormalization-group flow analysis of meson condensation in dense matter,” to appear.
- [12] G. Baym, *Phys. Rev. Lett.* **30** (1973) 1340.
- [13] For a powerful discussion, see H. Leutwyler, “On the foundations of chiral perturbation theory,” BUTP-93.24, *Ann. Phys.*, to be published.
- [14] For a review, see E. Jenkins and A.V. Manohar, in *Proc. of the Workshop on Effective Field Theories of the Standard Model*, Dobogókó, Hungary, Aug. 1991, ed. U.-G. Meissner (World Scientific, Singapore, 1992).
- [15] S. Weinberg, *Phys. Lett.* **B251** (1990) 288; *Nucl. Phys.* **B363** (1991) 3; *Phys. Lett.* **B295** (1992) 114.
- [16] T.-S. Park, D.-P. Min and M. Rho, *Phys. Repts.* **233** (1993) 341.
- [17] C.-H. Lee, H. Jung, D.-P. Min and M. Rho, *Phys. Lett.* **B326** (1993) 14.
- [18] C.-H. Lee, G.E. Brown and M. Rho, *Phys. Lett.* **B335** (1994) 266; C.-H. Lee, G.E. Brown, D.-P. Min and M. Rho, “An effective chiral Lagrangian approach to kaon-nuclear interactions: Kaonic atom and kaon condensation,” to appear in *Nucl. Phys.*.
- [19] E. Friedman, A. Gal and C.J. Batty, *Phys. Lett.* **B308** (1993) 6; *Nucl. Phys.* **A579** (1994) 518.
- [20] V. Thorsson, M. Prakash and J.M. Lattimer, *Nucl. Phys.* **A572** (1994) 693.
- [21] M. Prakash, T.L. Ainsworth and J.M. Lattimer, *Phys. Rev. Lett.* **61** (1988) 2518.